Author Index

Antonozzi, I.
——, Carducci, C., Vestri, L., Pontecorvi, A. and Moretti, F.
Rapid and sensitive method for high-performance liquid chromatographic analysis of pterins in biological fluids 319
Arrigoni Martelli, E., see Marzo, A. 313
Arzese, A., see Radin, L. 331
Badamchian, M.
——, Strickler, M. P., Stone, M. J. and Goldstein, A. L.
Rapid isolation of thymosin β_4 from thymosin fraction 5 by preparative high-performance liquid chromatography 291
Baig, M. S., see Qureshi, G. A. 237
Bear, G. R.
Universal detection and quantitation of surfactants by high-performance liquid chromatography by means of the evaporative light-scattering detector 91
Beebe, J. M.
——, Brown, P. R. and Turcotte, J. G.
Preparative-scale high-performance liquid chromatography of omega-3 polyunsaturated fatty acid esters derived from fish oil 369
Benedict, C. R., see Bodola, F. 281
Betto, P.
——, Ricciarello, G., Giambenedetti, M., Lucarelli, C., Ruggeri, S. and Stocchi, F.
Improved high-performance liquid chromatographic analysis with double detection system for L-DOPA, its metabolites and carbidopa in plasma of Parkinsonian patients under L-DOPA therapy 341
Biondi, P.A.
——, Manca, F., Negri, A., Secchi, C., Tedeschi, G. and Lucarelli, C.
Improved high-performance liquid chromatographic determination of bacterial collagenase activity in ointments 337
Bodola, F.
—— and Benedict, C. R.
Measurement of the release of adenine nucleotides during platelet aggregation by small-bore-column isocratic high-performance liquid chromatography 281
Bokx, P. K. de, see Schoenmakers, P. J. 201
Bornhop, D. J.
——, Schmidt, S. and Porter, N. L.
Use of two simultaneous detectors in capillary supercritical fluid chromatography 193
Botta, G. A., see Radin, L. 331
Brinkman, U. A. Th., see Kwakman, P. J. M. 139
Brown, P. R., see Beebe, J. M. 369
Brown, Y. L.
——, Kandrotas, R. J., Douglas, J. B. and Gal, P.
High-performance liquid chromatographic determination of indomethacin serum concentrations 275
Carducci, C., see Antonozzi, I. 319
Chan, Y.-P. M.
—— and Siu, T.-S. S.
Simultaneous quantitation of catecholamines and O-methylated metabolites in urine by isocratic ion-pairing high-performance liquid chromatography with amperometric detection 251
Chiodetti, L., see Knight, M. 361
Cooperman, B. S., see Ghrist, B. F. D. 1
Cordis, G. A.
——, Engelman, R. M. and Das, D. K.
Novel dual-wavelength monitoring approach for the improved rapid separation and estimation of adenine nucleotides and creatine phosphate by high-performance liquid chromatography 229
——, see Samanta, A. 221
Das, D. K., see Cordis, G. A. 229
——, see Samanta, A. 221
De Bokx, P. K., see Schoenmakers, P. J. 201
De Jong, G. J., see Kwakman, P. J. M. 139
Dewaele, C., see Nagels, L. J. 163
Douglas, J. B., see Brown, Y. L. 275
Edkins, T. J.
—— and Shelly, D. C.
Enhanced performance of a laser-induced fluorescence liquid chromatographic apparatus: a systems approach 109
Engelman, R. M., see Cordis, G. A. 229
Eriksson, B.-M.
—— and Persson, B.-A.
Determination of 5-hydroxytryptamine and 5-hydroxyindoleacetic acid in plasma by direct injection in coupled-column liquid chromatography with electrochemical detection 351
Evans, C. E.
——, Shabushnig, J. G. and McGuffin, V. L.
Experimental and theoretical model of refractive index artifacts in absorbance detection 119
Forsmo-Bruce, H., see Johansson, M. 301
Frei, R. W., see Kwakman, P. J. M. 139
Gal, P., see Brown, Y. L. 275

Geiser, F. O.
——, Yocklovich, S. G., Lurcott, S. M., Guthrie, J. W. and Levy, E. J.
Water as a stationary phase modifier in packed-column supercritical fluid chromatography. I. Separation of free fatty acids 173
Ghrist, B. F. D.
——, Cooperman, B. S. and Snyder, L. R.
Design of optimized high-performance liquid chromatographic gradients for the separation of either small or large molecules. I. Minimizing errors in computer simulations 1
—— and Snyder, L. R.
Design of optimized high-performance liquid chromatographic gradients for the separation of either small or large molecules. II. Background and theory 25
—— and Snyder, L. R.
Design of optimized high-performance liquid chromatographic gradients for the separation of either small or large molecules. III. An overall strategy and its application to several examples 43
Giambenedetti, M., see Betto, P. 341
Gluch, S., see Knight, M. 361
Goldstein, A. L., see Badamchian, M. 291
Grebow, P. E., see Hsu, S.-H. 215
Guthrie, J. W., see Geiser, F. O. 173
Haddad, P. R., see Sekulic, S. 65
—— and Sekulic, S.
Effects of peak tailing on computer optimisation procedures for high-performance liquid chromatography. II. An optimisation routine for tailed peaks 79
Hsu, S.-H.
——, Koerper, T. L., Tomlinson, B. J., Miksic, J. R. and Grebow, P. E.
Determination of a new inotropic agent in human plasma by high-performance liquid chromatography 215
Jacobsson, S., see Johansson, M. 301
Johansson, M.
——, Tufvesson Alm, A., Forsmo-Bruce, H., Jacobsson, S. and Westerlund, D.
Determination of noscapine and its metabolites in plasma by coupled-column liquid chromatography 301
Jong, G. J. de, see Kwakman, P. J. M. 139
Kamminga, D. A., see Kwakman, P. J. M. 139
Kandrotas, R. J., see Brown, Y. L. 275
Kanfer, I., see Skinner, M. 261
Kauffmann, J. M., see Nagels, L. J. 163
Knight, M.
——, Strickler, M. P., Stone, M. J., Chiodetti, L., Gluch, S. and Shinohara, T.
Scale-up methodology for the preparative purification of peptide M 361
Koerper, T. L., see Hsu, S.-H. 215
Krause, R. T.
—— and Wang, Y.
Oxidative detection of coulometrically reduced organonitro pesticides in reversed-phase high-performance liquid chromatography 151
Kwakman, P. J. M.
——, Mol, J. G. J., Kamminga, D. A., Frei, R. W., Brinkman, U. A. Th. and De Jong, G. J.
Rhodamine labelling reagent for the determination of chlorophenols by liquid chromatography with peroxyoxalate chemiluminescence detection 139
Levy, E. J., see Geiser, F. O. 173
Luber, Jr., J. M., see Rao, P. S. 269
Lucarelli, C., see Betto, P. 341
——, see Biondi, P. A. 337
——, see Marzo, A. 325
——, see Radin, L. 331
Lurcott, S. M., see Geiser, F. O. 173
McGuffin, V. L., see Evans, C. E. 119
Manca, F., see Biondi, P. A. 337
Martelli, E. Arrigoni, see Marzo, A. 313
Marzo, A.
——, Monti, N., Ripamonti, M. and Arrigoni Martelli, E.
Application of high-performance liquid chromatography to the analysis of propionyl-L-carnitine by a stereospecific enzyme assay 313
——, Reiner, A., Monti, N., Ripamonti, M. and Lucarelli, C.
High-performance liquid chromatographic evaluation of salicyloyl pyridixol and systemic metabolites in biological samples 325
Miksic, J. R., see Hsu, S.-H. 215
Mol, J. G. J., see Kwakman, P. J. M. 139
Monti, N., see Marzo, A. 313, 325
Moretti, F., see Antonozzi, I. 319
Nagels, L. J.
——, Kauffmann, J. M., Schuddinck, G., Dewaele, C., Patriarche, G. J. and Verzele, M.
Carbon–polymer chips as sensitive electrochemical detectors for micro-liquid chromatography 163
Negri, A., see Biondi, P. A. 337
Noguchi, K., see Seki, T. 245
Patriarche, G. J., see Nagels, L. J. 163
Persson, B.-A., see Eriksson, B.-M. 351
Pontecorvi, A., see Antonozzi, I. 319
Porter, N. L., see Bornhop, D. J. 193
Prasad, M. R., see Samanta, A. 221

Qureshi, G. A.
—— and Baig, M. S.
Quantitation of free amino acids in biological samples by high-performance liquid chromatography. Application of the method in evaluating amino acid levels in cerebrospinal fluid and plasma of patients with multiple sclerosis 237
Radin, L.
——, Arzese, A., Lucarelli, C. and Botta, G. A.
Rapid identification of *Bacteroides* species by high-performance liquid chromatography 331
Rao, P. S.
——, Rujikarn, N. and Luber, Jr., J. M.
High-performance liquid chromatographic method for the direct quantitation of oxy radicals in myocardium and blood by means of 1,3-dimethylthiourea and dimethyl sulfoxide 269
Reiner, A., see Marzo, A. 325
Ricciarello, G., see Betto, P. 341
Ripamonti, M., see Marzo, A. 313, 325
Ruggeri, S., see Betto, P. 341
Rujikarn, N., see Rao, P. S. 269
Samanta, A.
——, Cordis, G. A., Prasad, M. R. and Das, D. K.
Separation of fatty acid binding protein by high-performance mixed-mode chromatography 221
Schmidt, S., see Bornhop, D. J. 193
Schoenmakers, P. J.
——, Uunk, L. G. M. and De Bokx, P. K.
Effect of sample size on retention in packed column supercritical fluid chromatography. A method for characterizing stationary phase homogeneity 201
Schuddinck, G., see Nagels, L. J. 163
Secchi, C., see Biondi, P. A. 337
Seki, T.
——, Yanagihara, Y. and Noguchi, K.
Determination of salsolinol by ion-exchange chromatography with glycylglycine as the post-derivatizing agent 245
Sekulic, S., see Haddad, P. R. 79
—— and Haddad, P. R.
Effects of peak tailing on computer optimisation procedures for high-performance liquid chromatography. I. Characteristics of tailed peaks under optimisation conditions 65
Shabushnig, J. G., see Evans, C. E. 119
Shelly, D. C., see Edkins, T. J. 109
Shinohara, T., see Knight, M. 361
Siu, T.-S. S., see Chan, Y.-P. M. 251
Skinner, M.
—— and Kanfer, I.
High-performance liquid chromatographic analysis of josamycin in serum and urine 261
Smith, R. D., see Yonker, C. R. 183
Snyder, L. R., see Ghrist, B. F. D. 1, 25, 43
Stocchi, F., see Betto, P. 341
Stone, M. J., see Badamchian, M. 291
——, see Knight, M. 361
Strickler, M. P., see Badamchian, M. 291
——, see Knight, M. 361
Tedeschi, G., see Biondi, P. A. 337
Tomlinson, B. J., see Hsu, S.-H. 215
Tufvesson Alm, A., see Johansson, M. 301
Turcotte, J. G., see Beebe, J. M. 369
Uunk, L. G. M., see Schoenmakers, P. J. 201
Verzele, M., see Nagels, L. J. 163
Vestri, L., see Antonozzi, I. 319
Wang, Y., see Krause, R. T. 151
Westerlund, D., see Johansson, M. 301
Yanagihara, Y., see Seki, T. 245
Yocklovich, S. G., see Geiser, F. O. 173
Yonker, C. R.
—— and Smith, R. D.
Effect of the partial molar volume of the solute in the stationary phase on retention in supercritical fluid chromatography 183